

# COMPLEX NUMBER

## 1. DEFINITION

A number of the form  $a + ib$ , where  $a, b \in \mathbb{R}$  and  $i = \sqrt{-1}$ , is called a complex number and is denoted by 'Z'.

$$z = \boxed{a} + i\boxed{b}$$

$\downarrow$                      $\downarrow$   
 $\text{Re}(z)$             $\text{Im}(z)$

### 1.1 Conjugate of a Complex Number

For a given complex number  $z = a + ib$ , its conjugate ' $\bar{z}$ ' is defined as  $\bar{z} = a - ib$

## 2. ALGEBRA OF COMPLEX NUMBERS

Let  $z_1 = a + ib$  and  $z_2 = c + id$  be two complex numbers where  $a, b, c, d \in \mathbb{R}$  and  $i = \sqrt{-1}$ .

### 1. Addition :

$$\begin{aligned} z_1 + z_2 &= (a + bi) + (c + di) \\ &= (a + c) + (b + d) i \end{aligned}$$

### 2. Subtraction :

$$\begin{aligned} z_1 - z_2 &= (a + bi) - (c + di) \\ &= (a - c) + (b - d) i \end{aligned}$$

### 3. Multiplication :

$$\begin{aligned} z_1 \cdot z_2 &= (a + bi) (c + di) \\ &= a(c + di) + bi(c + di) \\ &= ac + adi + bci + bdi^2 \\ &= ac - bd + (ad + bc) i \end{aligned}$$

( $\because i^2 = -1$ )

### 4. Division :

$$\begin{aligned} \frac{z_1}{z_2} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \cdot \frac{c - di}{c - di} \\ &= \left( \frac{ac + bd}{c^2 + d^2} \right) + \left( \frac{bc - ad}{c^2 + d^2} \right) i \end{aligned}$$

Note...

1.  $a + ib = c + id$


$\Leftrightarrow a = c \ \& \ b = d$

2.  $i^{4k+r} = \begin{cases} 1; & r = 0 \\ i; & r = 1 \\ -1; & r = 2 \\ -i; & r = 3 \end{cases}$

3.  $\sqrt{\sqrt{\phantom{x}}} \sqrt{a} = \sqrt[4]{a}$  only if atleast one of either a or b is non-negative.

## 3. ARGAND PLANE


A complex number  $z = a + ib$  can be represented by a unique point P (a, b) in the argand plane.



$Z = a + ib$  is represented by a point P (a, b)

### 3.1 Modulus and Argument of Complex Number

If  $z = a + ib$  is a complex number



- (i) Distance of Z from origin is called as modulus of complex number Z.

It is denoted by  $r = |z| = \sqrt{a^2 + b^2}$

- (ii) Here,  $\theta$  i.e. angle made by OP with positive direction of real axis is called **argument of z**.



*Note...*

$z_1 > z_2$  or  $z_1 < z_2$  has no meaning but  $|z_1| > |z_2|$  or  $|z_1| < |z_2|$  holds meaning.

### 3.2 Principal Argument


The argument ' $\theta$ ' of complex number  $z = a + ib$  is called principal argument of z if  $-\pi < \theta \leq \pi$ .

Let  $\tan \alpha = \left| \frac{b}{a} \right|$ , and  $\theta$  be the arg (z).



In (iii) and (iv) principal argument is given by  $-\pi + \alpha$  and  $-\alpha$  respectively.

4. POLAR FORM



$$a = r \cos \theta \quad \& \quad b = r \sin \theta;$$

where  $r = |z|$  and  $\theta = \arg(z)$

$$\therefore z = a + ib$$

$$= r (\cos \theta + i \sin \theta)$$



Note...

$Z = re^{i\theta}$  is known as Euler's form; where  $r = |Z|$  &  $\theta = \arg(Z)$

5. SOME IMPORTANT PROPERTIES

1.  $\overline{(\overline{z})} = z$
2.  $z + \overline{z} = 2 \operatorname{Re}(z)$
3.  $z - \overline{z} = 2i \operatorname{Im}(z)$
4.  $\overline{z_1 + z_2} = \overline{z_1} + \overline{z_2}$
5.  $\overline{z_1 z_2} = \overline{z_1} \overline{z_2}$
6.  $|z| = 0 \Rightarrow z = 0$
7.  $z\overline{z} = |z|^2$
8.  $|z_1 z_2| = |z_1| |z_2|; \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$
9.  $|\overline{z}| = |z| = |-z|$
10.  $|z_1 \pm z_2|^2 = |z_1|^2 + |z_2|^2 \pm 2 \operatorname{Re}(z_1 \overline{z_2})$
11.  $|z_1 + z_2| \leq |z_1| + |z_2|$  (Triangle Inequality)
12.  $|z_1 - z_2| \geq ||z_1| - |z_2||$
13.  $|az_1 - bz_2|^2 + |bz_1 + az_2|^2 = (a^2 + b^2)(|z_1|^2 + |z_2|^2)$

$$14. \operatorname{amp}(z_1 \cdot z_2) = \operatorname{amp} z_1 + \operatorname{amp} z_2 + 2k\pi; k \in \mathbb{I}$$

$$15. \operatorname{amp} \left( \frac{y_0}{y_1} \right) = \operatorname{amp} z_1 - \operatorname{amp} z_2 + 2k\pi; k \in \mathbb{I}$$

$$16. \operatorname{amp}(z^n) = n \operatorname{amp}(z) + 2k\pi; k \in \mathbb{I}$$

6. DE-MOIVRE'S THEOREM

**Statement :**  $\cos n\theta + i \sin n\theta$  is the value or one of the values of  $(\cos \theta + i \sin \theta)^n$  according as if 'n' is integer or a rational number. The theorem is very useful in determining the roots of any complex quantity

7. CUBE ROOT OF UNITY

Roots of the equation  $x^3 = 1$  are called cube roots of unity.

$$x^3 - 1 = 0$$

$$(x - 1)(x^2 + x + 1) = 0$$

$$x = 1 \quad \text{or} \quad x^2 + x + 1 = 0$$

$$\text{i.e. } x = \frac{-1 + \sqrt{3}i}{2} \quad \text{or} \quad x = \frac{-1 - \sqrt{3}i}{2}$$

- (i) The cube roots of unity are  $1, \frac{-1 + i\sqrt{3}}{2}, \frac{-1 - i\sqrt{3}}{2}$ .
- (ii)  $W^3 = 1$
- (iii) If  $w$  is one of the imaginary cube roots of unity then  $1 + w + w^2 = 0$ .
- (iv) In general  $1 + w^r + w^{2r} = 0$ ; where  $r \in \mathbb{I}$  but is not the multiple of 3.
- (v) In polar form the cube roots of unity are :  
 $\cos 0 + i \sin 0; \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}, \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3}$
- (vi) The three cube roots of unity when plotted on the argand plane constitute the vertices of an equilateral triangle.
- (vii) The following factorisation should be remembered :

$$a^3 - b^3 = (a - b)(a - \omega b)(a - \omega^2 b);$$

$$x^2 + x + 1 = (x - \omega)(x - \omega^2);$$

$$a^3 + b^3 = (a + b)(a + \omega b)(a + \omega^2 b);$$

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a + \omega b + \omega^2 c)(a + \omega^2 b + \omega c)$$

### 8. 'n' n<sup>th</sup> ROOTS OF UNITY

Solution of equation  $x^n = 1$  is given by

$$x = \cos \frac{2k\pi}{n} + i \sin \frac{2k\pi}{n} ; k = 0, 1, 2, \dots, n-1$$

$$= e^{i\left(\frac{2k\pi}{n}\right)} ; k = 0, 1, \dots, n-1$$



Note...

- We may take any n consecutive integral values of k to get 'n' n<sup>th</sup> roots of unity.
- Sum of 'n' n<sup>th</sup> roots of unity is zero,  $n \in \mathbb{N}$
- The points represented by 'n' n<sup>th</sup> roots of unity are located at the vertices of regular polygon of n sides inscribed in a unit circle, centred at origin & one vertex being one +ve real axis.

#### Properties :

If  $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$  are the n, n<sup>th</sup> root of unity then :

- They are in G.P. with common ratio  $e^{i(2\pi/n)}$
- $1^p + \alpha_0^p + \alpha_1^p + \dots + \alpha_{n-1}^p = \begin{cases} 0, & \text{if } p \neq kn \\ n, & \text{if } p = kn \end{cases}$  where  $k \in \mathbb{Z}$
- $(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_{n-1}) = n$
- $(1 + \alpha_1)(1 + \alpha_2) \dots (1 + \alpha_{n-1}) = \begin{cases} 0, & \text{if } n \text{ is even} \\ 1, & \text{if } n \text{ is odd} \end{cases}$
- $1 \cdot \alpha_1 \cdot \alpha_2 \cdot \alpha_3 \dots \alpha_{n-1} = \begin{cases} -1, & \text{if } n \text{ is even} \\ 1, & \text{if } n \text{ is odd} \end{cases}$



Note...

- $\cos \theta + \cos 2\theta + \cos 3\theta + \dots + \cos n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \cos\left(\frac{n+1}{2}\theta\right)$
- $\sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \sin\left(\frac{n+1}{2}\theta\right)$

### 9. SQUARE ROOT OF COMPLEX NUMBER

Let  $x + iy = \sqrt{a + ib}$ , Squaring both sides, we get

$$(x + iy)^2 = a + ib$$

$$\text{i.e. } x^2 - y^2 = a, 2xy = b$$

Solving these equations, we get square roots of z.

### 10. LOCI IN COMPLEX PLANE

- $|z - z_0| = a$  represents circumference of circle, centred at  $z_0$ , radius a.
- $|z - z_0| < a$  represents interior of circle
- $|z - z_0| > a$  represents exterior of this circle.
- $|z - z_1| = |z - z_2|$  represents  $\perp$  bisector of segment with end points  $z_1$  &  $z_2$ .
- $\left| \frac{z - z_1}{z - z_2} \right| = k$  represents :  $\left\{ \begin{array}{l} \text{circle, } k \neq 1 \\ \perp \text{ bisector, } k = 1 \end{array} \right\}$
- $\arg(z) = \theta$  is a ray starting from origin (excluded) inclined at an  $\angle \theta$  with real axis.
- Circle described on line segment joining  $z_1$  &  $z_2$  as diameter is :  
 $(z - z_1)(\bar{z} - \bar{z}_2) + (z - z_2)(\bar{z} - \bar{z}_1) = 0$ .
- Four pts.  $z_1, z_2, z_3, z_4$  in anticlockwise order will be concyclic, if & only if

$$\theta = \arg\left(\frac{z_2 - z_4}{z_1 - z_4}\right) = \arg\left(\frac{z_2 - z_3}{z_1 - z_3}\right)$$

$$\Rightarrow \arg\left(\frac{z_2 - z_4}{z_1 - z_4}\right) - \arg\left(\frac{z_2 - z_3}{z_1 - z_3}\right) = 2n\pi ; (n \in \mathbb{I})$$


$$\Rightarrow \arg\left[\left(\frac{z_2 - z_4}{z_1 - z_4}\right)\left(\frac{z_1 - z_3}{z_2 - z_3}\right)\right] = 2n\pi$$

$$\Rightarrow \left(\frac{z_2 - z_4}{z_1 - z_4}\right) \times \left(\frac{z_1 - z_3}{z_2 - z_3}\right) \text{ is real \& positive.}$$

**11. VECTORIAL REPRESENTATION OF A COMPLEX**

Every complex number can be considered as if it is the position vector of that point. If the point P represents the complex number z then,

$$\vec{OP} = z \quad \& \quad |\vec{OP}| = |z|.$$



**12. SOME IMPORTANT RESULTS**

- (i) If  $z_1$  and  $z_2$  are two complex numbers, then the distance between  $z_1$  and  $z_2$  is  $|z_2 - z_1|$ .
- (ii) Segment joining points A ( $z_1$ ) and B( $z_2$ ) is divided by point P ( $z$ ) in the ratio  $m_1 : m_2$

then  $z = \frac{m_1 z_2 + m_2 z_1}{m_1 + m_2}$ ,  $m_1$  and  $m_2$  are real.

- (iii) The equation of the line joining  $z_1$  and  $z_2$  is given by

$$\begin{vmatrix} z & \bar{z} \\ z_1 & \bar{z}_1 \\ z_2 & \bar{z}_2 \end{vmatrix} = 0 \quad (\text{non parametric form})$$

Or

$$\frac{z - z_1}{\bar{z} - \bar{z}_1} = \frac{z - z_2}{\bar{z} - \bar{z}_2}$$

- (iv)  $\bar{a}z + a\bar{z} + b = 0$  represents general form of line.
- (v) The general eqn. of circle is :

$$z\bar{z} + a\bar{z} + \bar{a}z + b = 0 \quad (\text{where } b \text{ is real no.})$$

Centre :  $(-a)$  & radius  $\sqrt{|a|^2 - b} = \sqrt{a\bar{a} - b}$ .

- (vi) Circle described on line segment joining  $z_1$  &  $z_2$  as diameter is :

$$(z - z_1)(\bar{z} - \bar{z}_2) + (z - z_2)(\bar{z} - \bar{z}_1) = 0.$$

*Note...*

- (i) If  $\vec{OP} = z = r e^{i\theta}$  then  $\vec{OQ} = z_1 = r e^{i(\theta + \phi)} = z \cdot e^{i\phi}$ .

If  $\vec{OP}$  and  $\vec{OQ}$  are of unequal magnitude then

$$\hat{OQ} = \hat{OP} e^{i\phi}$$

- (ii) If  $z_1, z_2, z_3$ , are three vertices of a triangle ABC described in the counter-clock wise sense, then

$$\frac{z_3 - z_1}{z_2 - z_1} = \frac{AC}{AB} (\cos \alpha + i \sin \alpha) = \frac{AC}{AB} \cdot e^{i\alpha} = \frac{|z_3 - z_1|}{|z_2 - z_1|} \cdot e^{i\alpha}$$

(vii) Four pts.  $z_1, z_2, z_3, z_4$  in anticlockwise order will be concyclic, if & only if

$$\theta = \arg\left(\frac{z_2 - z_4}{z_1 - z_4}\right) = \arg\left(\frac{z_2 - z_3}{z_1 - z_3}\right)$$

$$\Rightarrow \arg\left(\frac{z_2 - z_4}{z_1 - z_4}\right) - \arg\left(\frac{z_2 - z_3}{z_1 - z_3}\right) = 2n\pi; (n \in \mathbb{I})$$

$$\Rightarrow \arg\left[\left(\frac{z_2 - z_4}{z_1 - z_4}\right)\left(\frac{z_1 - z_3}{z_2 - z_3}\right)\right] = 2n\pi$$

$$\Rightarrow \left(\frac{z_2 - z_4}{z_1 - z_4}\right) \times \left(\frac{z_1 - z_3}{z_2 - z_3}\right) \text{ is real \& positive.}$$

(viii) If  $z_1, z_2, z_3$  are the vertices of an equilateral triangle where  $z_0$  is its circumcentre then

$$(a) \frac{1}{z_2 - z_3} + \frac{1}{z_3 - z_1} + \frac{1}{z_1 - z_2} = 0$$

$$(b) z_0^1 + z_1^1 + z_2^1 - z_1 z_2 - z_2 z_3 - z_3 z_1 = 0$$

$$(c) z_0^1 + z_1^1 + z_2^1 = 3 z_0^1$$

(ix) If A, B, C & D are four points representing the complex numbers  $z_1, z_2, z_3$  &  $z_4$  then

$$AB \parallel CD \text{ if } \frac{z_4 - z_3}{z_2 - z_1} \text{ is purely real ;}$$

$$AB \perp CD \text{ if } \frac{z_4 - z_3}{z_2 - z_1} \text{ is purely imaginary ]}$$

(x) Two points P ( $z_1$ ) and Q ( $z_2$ ) lie on the same side or opposite side of the line  $\bar{a}z + a\bar{z} + b$  accordingly as  $\bar{a}z_1 + a\bar{z}_1 + b$  and  $\bar{a}z_2 + a\bar{z}_2 + b$  have same sign or opposite sign.

**Important Identities**

$$(i) x^2 + x + 1 = (x - \omega)(x - \omega^2)$$

$$(ii) x^2 - x + 1 = (x + \omega)(x + \omega^2)$$

$$(iii) x^2 + xy + y^2 = (x - y\omega)(x - y\omega^2)$$

$$(iv) x^2 - xy + y^2 = (x + y\omega)(x + y\omega^2)$$

$$(v) x^2 + y^2 = (x + iy)(x - iy)$$

$$(vi) x^3 + y^3 = (x + y)(x + y\omega)(x + y\omega^2)$$

$$(vii) x^3 - y^3 = (x - y)(x - y\omega)(x - y\omega^2)$$

$$(viii) x^2 + y^2 + z^2 - xy - yz - zx = (x + y\omega + z\omega^2)(x + y\omega^2 + z\omega)$$

$$\text{or } (x\omega + y\omega^2 + z)(x\omega^2 + y\omega + z)$$

$$\text{or } (x\omega + y + z\omega^2)(x\omega^2 + y + z\omega)$$

$$(ix) x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x + \omega y + \omega^2 z)(x + \omega^2 y + \omega z)$$

# QUADRATIC EQUATION

## 1. QUADRATIC EXPRESSION

The general form of a quadratic expression in  $x$  is,

$$f(x) = ax^2 + bx + c, \text{ where } a, b, c \in \mathbb{R} \text{ \& } a \neq 0.$$

and general form of a quadratic equation in  $x$  is,

$$ax^2 + bx + c = 0, \text{ where } a, b, c \in \mathbb{R} \text{ \& } a \neq 0.$$

## 2. ROOTS OF QUADRATIC EQUATION

(a) **The solution of the quadratic equation,**

$$ax^2 + bx + c = 0 \text{ is given by } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

The expression  $D = b^2 - 4ac$  is called the discriminant of the quadratic equation.

(b) **If  $\alpha$  &  $\beta$  are the roots of the quadratic equation**

$$ax^2 + bx + c = 0, \text{ then ;}$$

$$(i) \alpha + \beta = -b/a \quad (ii) \alpha \beta = c/a$$

$$(iii) |\alpha - \beta| = \frac{\sqrt{D}}{|a|}.$$

(c) **A quadratic equation whose roots are  $\alpha$  &  $\beta$  is**  
 $(x - \alpha)(x - \beta) = 0$  i.e.

$$x^2 - (\alpha + \beta)x + \alpha\beta = 0 \quad \text{i.e.}$$

$$x^2 - (\text{sum of roots})x + \text{product of roots} = 0.$$

*Note...* 

$$y = (ax^2 + bx + c) \equiv a(x - \alpha)(x - \beta)$$

$$= a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a}$$

## 3. NATURE OF ROOTS

(a) **Consider the quadratic equation  $ax^2 + bx + c = 0$  where  $a, b, c \in \mathbb{R}$  &  $a \neq 0$  then;**

(i)  $D > 0 \Leftrightarrow$  roots are real & distinct (unequal).

(ii)  $D = 0 \Leftrightarrow$  roots are real & coincident (equal).

(iii)  $D < 0 \Leftrightarrow$  roots are imaginary.

(iv) If  $p + iq$  is one root of a quadratic equation, then the other must be the conjugate  $p - iq$  & vice versa. ( $p, q \in \mathbb{R}$  &  $i = \sqrt{-1}$ ).

(b) **Consider the quadratic equation  $ax^2 + bx + c = 0$  where  $a, b, c \in \mathbb{Q}$  &  $a \neq 0$  then;**

(i) If  $D > 0$  & is a perfect square, then roots are rational & unequal.

(ii) If  $\alpha = p + \sqrt{q}$  is one root in this case, (where  $p$  is rational &  $\sqrt{q}$  is a surd) then the other root must be the conjugate of it i.e.  $\beta = p - \sqrt{q}$  & vice versa.

*Note...* 

Remember that a quadratic equation cannot have three different roots & if it has, it becomes an identity.

**4. GRAPH OF QUADRATIC EXPRESSION**

Consider the quadratic expression,  $y = ax^2 + bx + c$ ,  $a \neq 0$  &  $a, b, c \in \mathbb{R}$  then ;

- (i) The graph between  $x, y$  is always a parabola. If  $a > 0$  then the shape of the parabola is concave upwards & if  $a < 0$  then the shape of the parabola is concave downwards.
- (ii)  $y > 0 \forall x \in \mathbb{R}$ , only if  $a > 0$  &  $D < 0$
- (iii)  $y < 0 \forall x \in \mathbb{R}$ , only if  $a < 0$  &  $D < 0$


**5. SOLUTION OF QUADRATIC INEQUALITIES**

$ax^2 + bx + c > 0$  ( $a \neq 0$ ).

- (i) If  $D > 0$ , then the equation  $ax^2 + bx + c = 0$  has two different roots ( $x_1 < x_2$ ).

Then  $a > 0 \Rightarrow x \in (-\infty, x_1) \cup (x_2, \infty)$

$a < 0 \Rightarrow x \in (x_1, x_2)$




- (ii) Inequalities of the form  $\frac{P(x)}{Q(x)} \geq 0$  can be quickly solved using the method of intervals (wavy curve).

**6. MAX. & MIN. VALUE OF QUADRATIC EXPRESSION**

Maximum & Minimum Value of  $y = ax^2 + bx + c$  occurs at  $x = -(b/2a)$  according as :

**For  $a > 0$ , we have :**


$$y \in \left[ \frac{4ac - b^2}{4a}, \infty \right)$$



$$y_{\min} = \frac{-D}{4a} \text{ at } x = \frac{-b}{2a}, \text{ and } y_{\max} \rightarrow \infty$$

**For  $a < 0$ , we have :**

$$y \in \left( -\infty, \frac{4ac - b^2}{4a} \right]$$



$$y_{\max} = \frac{-D}{4a} \text{ at } x = \frac{-b}{2a}, \text{ and } y_{\min} \rightarrow -\infty$$



## 7. THEORY OF EQUATIONS

If  $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$  are the roots of the  $n^{\text{th}}$  degree polynomial equation :

$$f(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n = 0$$

where  $a_0, a_1, \dots, a_n$  are all real &  $a_0 \neq 0$ ,

Then,

$$\sum \alpha_1 = -\frac{a_1}{a_0};$$

$$\sum \alpha_1 \alpha_2 = \frac{a_2}{a_0};$$

$$\sum \alpha_1 \alpha_2 \alpha_3 = -\frac{a_3}{a_0};$$

.....

$$\alpha_1 \alpha_2 \alpha_3 \dots \alpha_n = (-1)^n \frac{a_n}{a_0}$$

## 8. LOCATION OF ROOTS

Let  $f(x) = ax^2 + bx + c$ , where  $a > 0$  &  $a, b, c \in \mathbb{R}$ .

- (i) Conditions for both the roots of  $f(x) = 0$  to be greater than a specified number 'k' are :  
 $D \geq 0$  &  $f(k) > 0$  &  $(-b/2a) > k$ .
- (ii) Conditions for both roots of  $f(x) = 0$  to lie on either side of the number 'k' (in other words the number 'k' lies between the roots of  $f(x) = 0$  is :  
 $a f(k) < 0$ .
- (iii) Conditions for exactly one root of  $f(x) = 0$  to lie in the interval  $(k_1, k_2)$  i.e.  $k_1 < x < k_2$  are :  
 $D > 0$  &  $f(k_1) \cdot f(k_2) < 0$ .
- (iv) Conditions that both roots of  $f(x) = 0$  to be confined between the numbers  $k_1$  &  $k_2$  are  $(k_1 < k_2)$  :  
 $D \geq 0$  &  $f(k_1) > 0$  &  $f(k_2) > 0$  &  $k_1 < (-b/2a) < k_2$ .

Note...

**Remainder Theorem :** If  $f(x)$  is a polynomial, then  $f(h)$  is the remainder when  $f(x)$  is divided by  $x - h$ .

**Factor theorem :** If  $x = h$  is a root of equation  $f(x) = 0$ , then  $x-h$  is a factor of  $f(x)$  and conversely.

## 9. MAX. & MIN. VALUES OF RATIONAL EXPRESSION

Here we shall find the values attained by a rational

expression of the form  $\frac{a_1x^2 + b_1x + c_1}{a_2x^2 + b_2x + c_2}$  for real values

of  $x$ .

**Example No. 4** will make the method clear.

## 10. COMMON ROOTS

### (a) Only One Common Root

Let  $\alpha$  be the common root of  $ax^2 + bx + c = 0$  &  $a'x^2 + b'x + c' = 0$ , such that  $a, a' \neq 0$  and  $a' \neq a$ .

Then, the condition for one common root is :

$$(ca' - c'a)^2 = (ab' - a'b)(bc' - b'c).$$

### (b) Two Common Roots

Let  $\alpha, \beta$  be the two common roots of

$$ax^2 + bx + c = 0 \text{ \& \ } a'x^2 + b'x + c' = 0,$$

such that  $a, a' \neq 0$ .

Then, the condition for two common roots is :

$$\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$$

### 11. RESOLUTION INTO TWO LINEAR FACTORS

The condition that a quadratic function

$$f(x, y) = ax^2 + 2hxy + by^2 + 2gx + 2fy + c$$

may be resolved into two linear factors is that ;

$$abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

OR 
$$\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0$$

### 12. FORMATION OF A POLYNOMIAL EQUATION

If  $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$  are the roots of the  $n^{\text{th}}$  degree polynomial equation, then the equation is

$$x^n - S_1x^{n-1} + S_2x^{n-2} + S_3x^{n-3} + \dots + (-1)^n S_n = 0$$

where  $S_k$  denotes the sum of the products of roots taken  $k$  at a time.

#### Particular Cases

(a) **Quadratic Equation** if  $\alpha, \beta$  be the roots the quadratic equation, then the equation is :

$$x^2 - S_1x + S_2 = 0 \quad \text{i.e.} \quad x^2 - (\alpha + \beta)x + \alpha\beta = 0$$

(b) **Cubic Equation** if  $\alpha, \beta, \gamma$  be the roots the cubic equation, then the equation is :

$$x^3 - S_1x^2 + S_2x - S_3 = 0 \quad \text{i.e.}$$

$$x^3 - (\alpha + \beta + \gamma)x^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)x - \alpha\beta\gamma = 0$$

(i) If  $\alpha$  is a root of equation  $f(x) = 0$ , the polynomial  $f(x)$  is exactly divisible by  $(x - \alpha)$ . In other words,  $(x - \alpha)$  is a factor of  $f(x)$  and conversely.

(ii) Every equation of  $n^{\text{th}}$  degree ( $n \geq 1$ ) has exactly  $n$  roots & if the equation has more than  $n$  roots, it is an identity.

(iii) If there be any two real numbers 'a' & 'b' such that  $f(a)$  &  $f(b)$  are of opposite signs, then  $f(x) = 0$  must have atleast one real root between 'a' and 'b'.

(iv) Every equation  $f(x) = 0$  of degree odd has atleast one real root of a sign opposite to that of its last term.

### 13. TRANSFORMATION OF EQUATIONS

(i) To obtain an equation whose roots are reciprocals of the roots of a given equation, it is obtained by replacing  $x$  by  $1/x$  in the given equation

(ii) Transformation of an equation to another equation whose roots are negative of the roots of a given equation—replace  $x$  by  $-x$ .

(iii) Transformation of an equation to another equation whose roots are square of the roots of a given equation—replace  $x$  by  $\sqrt{x}$ .

(iv) Transformation of an equation to another equation whose roots are cubes of the roots of a given equation—replace  $x$  by  $x^{1/3}$ .